



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 241/08, 241/04, 243/08, 401/12, 409/06, A61K 31/495		A1	(11) International Publication Number: WO 97/40023
			(43) International Publication Date: 30 October 1997 (30.10.97)

<p>(21) International Application Number: PCT/DK97/00186</p> <p>(22) International Filing Date: 24 April 1997 (24.04.97)</p> <p>(30) Priority Data: 0489/96 24 April 1996 (24.04.96) DK 1344/96 26 November 1996 (26.11.96) DK</p> <p>(71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): HANSEN, Thomas, Kruse [DK/DK]; 78 Tibbevangen, DK-2730 Herlev (DK). PESCHKE, Bernd [DE/DK]; 56 Eskebjerggaard, DK-2760 Maaloev (DK). ANDERSEN, Knud, Erik [DK/DK]; 122 Nøddelunden, DK-2765 Smørum (DK).</p> <p>(74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsvaerd (DK).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
--	---

(54) Title: COMPOUNDS WITH GROWTH HORMONE RELEASING PROPERTIES

(57) Abstract

The present invention relates to novel compounds of general formula (I) which can be used for treating medical disorders resulting from a deficiency in growth hormone.

(I)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOUNDS WITH GROWTH HORMONE RELEASING PROPERTIES

FIELD OF INVENTION

- 5 The present invention relates to novel compounds, compositions containing them, and their use for treating medical disorders resulting from a deficiency in growth hormone.

BACKGROUND OF THE INVENTION

10

Growth hormone is a hormone which stimulates growth of all tissues capable of growing. In addition, growth hormone is known to have a number of effects on metabolic processes, e.g., stimulation of protein synthesis and free fatty acid mobilization and to cause a switch in energy metabolism from carbohydrate to
15 fatty acid metabolism. Deficiency in growth hormone can result in a number of severe medical disorders, e.g., dwarfism.

Growth hormone is released from the pituitary. The release is under tight control of a number of hormones and neurotransmitters either directly or indirectly.
20 Growth hormone release can be stimulated by growth hormone releasing hormone (GHRH) and inhibited by somatostatin. In both cases the hormones are released from the hypothalamus but their action is mediated primarily via specific receptors located in the pituitary. Other compounds which stimulate the release of growth hormone from the pituitary have also been described. For example
25 arginine, L-3,4-dihydroxyphenylalanine (L-Dopa), glucagon, vasopressin, PACAP (pituitary adenylyl cyclase activating peptide), muscarinic receptor agonists and a synthetic hexapeptide, GHRP (growth hormone releasing peptide) release endogenous growth hormone either by a direct effect on the pituitary or by affecting the release of GHRH and/or somatostatin from the hypothalamus.

30

In disorders or conditions where increased levels of growth hormone is desired, the protein nature of growth hormone makes anything but parenteral administration non-viable. Furthermore, other directly acting natural secretagogues, e.g., GHRH and PACAP, are longer polypeptides for which
35 reason oral administration of them is not viable.

The use of certain compounds for increasing the levels of growth hormone in mammals has previously been proposed, e.g. in EP 18 072, EP 83 864, WO 89/07110, WO 89/01711, WO 89/10933, WO 88/9780, WO 83/02272, WO 91/18016, WO 92/01711, WO 93/04081, WO 95/17422, WO 95/17423 and WO 95/14666.

The composition of growth hormone releasing compounds is important for their growth hormone releasing potency as well as their bioavailability. It is therefore an object of the present invention to provide novel compounds with growth hormone releasing properties which have improved properties relative to known compounds of this type.


SUMMARY OF THE INVENTION

In accordance with the present invention there is provided compounds which act directly on the pituitary cells under normal experimental conditions in vitro to release growth hormone therefrom.

These growth hormone releasing compounds can be utilized in vitro as unique research tools for understanding, inter alia, how growth hormone secretion is regulated at the pituitary level.

Moreover, the growth hormone releasing compounds of the present invention can also be administered in vivo to increase growth hormone release.

Accordingly, the present invention relates to a compound of general formula I



formula I

wherein

R^1 is hydrogen, or C_{1-6} -alkyl optionally substituted with aryl;

5 a and b are independently 1 or 2;

c and d are independently 0, 1, or 2;

c + d is 0, 1, or 2;

10


$R^2, R^3, R^4, R^6, R^7, R^8, R^9, R^{10}$, and R^{11} are independently hydrogen, aryl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$, or C_{1-6} -alkyl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$;

15


R^3 and R^4 can be taken together to form $=O$ or $=S$;

R^8 and R^9 can be taken together to form $=O$ or $=S$;

20 R^5 is hydrogen,



or C_{1-6} -alkyl optionally substituted with aryl, hydroxy, C_{3-6} -cycloalkyl, amino, -





25

R, R' and R'' are independently hydrogen or C_{1-6} -alkyl;


R^{30} and R^{31} are independently C_{1-6} -alkyl optionally substituted with aryl, hydroxy, C_{3-6} -cycloalkyl, or amino;

30

R^{22} , R^{23} , R^{24} , R^{25} , R^{26} , and R^{27} are independently hydrogen or C_{1-6} -alkyl;
when R^3 and R^4 are taken together to form $=O$ or $=S$, E is



- 5 when R^3 or R^4 is hydrogen, aryl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$, or C_{1-6} -alkyl, optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$, E is



- 10 or $R^{17} \cdot NH - (CR^{18}R^{19})_p - (CH_2)_m - Q - (CHR^{20})_o - (CH_2)_n -$

R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

- any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;
- 15

n, m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;


5

M is $-\text{CH}=\text{CR}^{21}-$, $-\text{O}-$, $-\text{S}-$, or a valence bond ;

Q is $-\text{CH}=\text{CR}^{21}-$, $-\text{O}-$, or $-\text{S}-$;

10 R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

G is hydrogen,




wherein R^{28} is hydrogen, halogen, C_{1-6} -alkyl, C_{1-6} -alkoxy or aryl;

r is 0, 1, or 2;

20

J is hydrogen,



wherein R^{29} is hydrogen, halogen, C_{1-6} -alkyl, C_{1-6} -alkoxy or aryl;
t is 0, 1, or 2;

- 5 or a pharmaceutically acceptable salt thereof.

The compounds of formula I comprise any optical isomers thereof, in the form of separated, pure or partially purified optical isomers or racemic mixtures thereof.

- 10 The compounds of formula II-V are independently preferred embodiments of the compound of formula I, thus any reference in the present description to formula I also means reference to formula II-V.

- In one embodiment of the compound of the above formula I R^1 is preferably C_{1-6} -alkyl, more preferred C_{1-4} -alkyl, such as methyl.
15

In a second embodiment of the compound of the above formula I a is preferably 1.

- In a further embodiment of the compound of the above formula I b is preferably 1.
20

In a still further embodiment of the compound of the above formula I c is preferably 0 or 1.

- In a further embodiment of the compound of the above formula I d is preferably 0 or 1.
25

In a still further embodiment of the compound of the above formula I c+d is preferably 1.

- 30 In a further embodiment of the compound of the above formula I R^2 is preferably hydrogen or C_{1-4} -alkyl, more preferred hydrogen.

In a still further embodiment of the compound of the above formula I R^3 and R^4 are preferably taken together to form =O or =S, more preferred =O.


In another embodiment of the compound of the above formula I R^3 and R^4 are independently of each other hydrogen.

In a further embodiment of the compound of the above formula I R^5 is preferably
 5 hydrogen, C_{1-6} -alkylsulfonyl, such as methylsulfonyl, or C_{1-6} -alkyl, more preferred C_{1-4} -alkyl, such as methyl.

In a still further embodiment of the compound of the above formula I R^6 , R^7 , R^8
 10 and R^9 are independently of each other preferably hydrogen or C_{1-6} -alkyl, more preferred C_{1-4} -alkyl, such as methyl.

In a further embodiment of the compound of the above formula I R^{10} and R^{11} are
 15 independently of each other preferably hydrogen or C_{1-4} -alkyl, more preferred hydrogen.

In a still further embodiment of the compound of the above formula I E is preferably



20

wherein R^{17} is hydrogen or C_{1-4} alkyl, preferably methyl,

R^{18} is hydrogen or C_{1-4} alkyl, preferably methyl, and


R^{19} is hydrogen or C_{1-4} alkyl, preferably methyl, or

R^{18} and R^{19} are joined together to form a C_{1-6} -alkylene bridge, preferably a
 25 C_{1-4} -alkylene bridge, such as a trimethylene-bridge, and

n is 0 or 1, preferably 0,

p is 1, m is 1 and M is $-CH=CR^{21}$, wherein R^{21} is hydrogen or C_{1-4} alkyl,
 preferably methyl,

30 more preferred E is $-CH=CH-CH_2-C(CH_3)_2NH_2$, $-CH=C(CH_3)-CH_2-C(CH_3)_2NH_2$, $-CH=CH-CH_2-C(CH_3)_2-NH(CH_3)$ or




In a further embodiment of the compound of the above formula I G is preferably



wherein R^{28} is hydrogen or aryl, preferably phenyl,
more preferred G is 2-naphthyl or biphenyl-4-yl.

5

In a still further embodiment of the compound of the above formula I J is preferably




wherein R^{29} is hydrogen or C_{1-6} -alkyl, preferably hydrogen,
more preferred J is phenyl or 2-thienyl.

10

Preferred compounds of the invention are:


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:

15




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-(cyclopropylmethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:

20




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-(hydroxymethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:

5




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxo-1,4-diazepane-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methyl amide:

10




Piperidine-4-carboxylic acid N-methyl-N-((1R)-2-(2-naphthyl)-1-((2R)-2-((2-naphthyl)methyl)-3-oxopiperazine-1-carbonyl)ethyl) amide:




5

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-(dimethyl-carbamoylmethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methyl amide:




10

(2E)-5-Methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:




(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R,6S)-2-benzyl-6-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:




5

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(biphenyl-4-yl)ethyl)-N-methylamide:




10


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-methanesulfonylpiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide:




5 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-methanesulfonylpiperazine-1-carbonyl)-2-benzyloxyethyl)-N-methylamide:




10 (2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.




(2E)-5-Methyl-5-methylaminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.




5 (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide



10 (2E)-5-Methyl-5-aminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxo piperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-methyl-N-((1R)-2-(2-naphthyl)-1-((2R)-3-oxo-2-((2-thienyl)methyl)piperazine-1-carbonyl)ethyl)amide




(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-2-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide

5




(2E) 5-Methyl-5-methylamino-hex-2-enoic acid N-methyl-N-((1R)-1-(2-naphthyl)methyl-2-oxo-2-((2R)-3-oxo-2-((2-thienyl)methyl)piperazin-1-yl)ethyl)amide.

10



(2E)-5-Amino-5-methyl-hex-2-enoic acid (2-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-(1R)-1-(naphth-2-ylmethyl)-2-oxo-ethyl)-methyl-amide.

15



It is believed that compounds of formula I exhibit an improved resistance to proteolytic degradation by enzymes compared to that of the peptides suggested in the prior literature, due to the lack of natural peptide bonds. The increased resistance to proteolytic degradation combined with the reduced size of the compounds of the invention in comparison with known growth hormone releasing peptides is expected to improve their bioavailability compared to that of the peptides suggested in the prior literature.

In the above structural formulas and throughout the present specification, the following terms have the indicated meanings:

The C₁₋₆-alkyl, or C₁₋₄-alkyl groups specified above are intended to include those alkyl groups of the designated length in either a linear or branched or cyclic configuration. Examples of linear alkyl are methyl, ethyl, propyl, butyl, pentyl, and hexyl. Examples of branched alkyl are isopropyl, sec-butyl, tert-butyl, isopentyl, and isohexyl. Examples of cyclic alkyl are C₃₋₆-cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The C₁₋₆-alkoxy groups specified above are intended to include those alkoxy groups of the designated length in either a linear or branched or cyclic configuration. Examples of linear alkoxy are methoxy, ethoxy, propoxy, butoxy, pentoxy, and hexoxy. Examples of branched alkoxy are isopropoxy, sec-butoxy, tert-butoxy, isopentoxy, and isohexoxy. Examples of cyclic alkoxy are cyclopropyloxy, cyclobutyloxy, cyclopentyloxy and cyclohexyloxy.

In the present context, the term "aryl" is intended to include aromatic rings, such as carbocyclic and heterocyclic aromatic rings selected from the group consisting of phenyl, naphthyl, pyridyl, 1-H-tetrazol-5-yl, thiazolyl, imidazolyl, indolyl, pyrimidinyl, thiadiazolyl, pyrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiophenyl,

quinolinyl, pyrazinyl, or isothiazolyl, optionally substituted by one or more C₁₋₆-alkyl, C₁₋₆-alkoxy, halogen, amino or aryl. Aryl is preferably phenyl, thienyl, imidazolyl, oxadiazolyl, pyridyl, indolyl, quinolinyl or naphthyl optionally substituted with halogen, amino, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy.

5

The term "halogen" is intended to include chlorine (Cl), fluorine (F), bromine (Br) and iodine (I).

10 The compounds of the present invention may have one or more asymmetric centres and it is intended that stereoisomers, as separated, pure or partially purified stereoisomers or racemic mixtures thereof are included in the scope of the invention.

15 The compounds of the present invention may optionally be on a pharmaceutically acceptable salt form such as the pharmaceutically acceptable acid addition salts of compounds of formula I which include those prepared by reacting the compound of formula I with an inorganic or organic acid such as hydrochloric, hydrobromic, sulfuric, acetic, phosphoric, lactic, maleic, phthalic, citric, glutaric, gluconic, methanesulfonic, salicylic, succinic, tartaric, toluenesulfonic, 20 trifluoroacetic, sulfamic or fumaric acid.

The compounds of formula I may be administered in pharmaceutically acceptable acid addition salt form or, where appropriate, as a alkali metal or alkaline earth metal or lower alkylammonium salt. Such salt forms are believed to exhibit 25 approximately the same order of activity as the free base forms.

In another aspect, the present invention relates to a pharmaceutical composition comprising, as an active ingredient, a compound of the general formula I or a pharmaceutically acceptable salt thereof together with a pharmaceutically 30 acceptable carrier or diluent.

Pharmaceutical compositions containing a compound of the present invention may be prepared by conventional techniques, e.g. as described in Remington's Pharmaceutical Sciences, 1985 or in Remington: The Science and Practice of 35 Pharmacy, 19th Edition (1995). The compositions may appear in conventional

forms, for example capsules, tablets, aerosols, solutions, suspensions or topical applications.

5 The pharmaceutical carrier or diluent employed may be a conventional solid or liquid carrier. Examples of solid carriers are lactose, terra alba, sucrose, cyclodextrin, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid or lower alkyl ethers of cellulose. Examples of liquid carriers are syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene or water.

10 Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

15 If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

20

A typical tablet which may be prepared by conventional tableting techniques may contain:

Core:

	Active compound (as free compound or salt thereof)	100mg
	Colloidal silicon dioxide (Aerosil)	1.5mg
	Cellulose, microcryst. (Avicel)	70mg
5	Modified cellulose gum (Ac-Di-Sol)	7.5mg
	Magnesium stearate	

Coating:

	HPMC approx.	9mg
10	*Mywacett 9-40 T approx.	0.9mg

*Acylated monoglyceride used as plasticizer for film coating.

15 For nasal administration, the preparation may contain a compound of formula I dissolved or suspended in a liquid carrier, in particular an aqueous carrier, for aerosol application. The carrier may contain additives such as solubilizing agents, e.g. propylene glycol, surfactants, absorption enhancers such as lecithin (phosphatidylcholine) or cyclodextrin, or preservatives such as parabenes.

20 Generally, the compounds of the present invention are dispensed in unit dosage form comprising 50-200 mg of active ingredient together with a pharmaceutically acceptable carrier per unit dosage.

25 The dosage of the compounds according to this invention is suitably 0.1-500 mg/day, e.g. from about 5 to about 50 mg, such as about 10 mg per dose, when administered to patients, e.g. humans, as a drug.

30 It has been demonstrated that compounds of the general formula I possess the ability to release endogenous growth hormone *in vivo*. The compounds may therefore be used in the treatment of conditions which require increased plasma growth hormone levels such as in growth hormone deficient humans or in elderly patients or livestock.

35 Thus, in a particular aspect, the present invention relates to a pharmaceutical composition for stimulating the release of growth hormone from the pituitary, the

composition comprising, as an active ingredient, a compound of the general formula I or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

5 In a further aspect, the present invention relates to a method of stimulating the release of growth hormone from the pituitary, the method comprising administering to a subject in need thereof an effective amount of a compound of the general formula I or a pharmaceutically acceptable salt thereof.

10 In a still further aspect, the present invention relates to the use of a compound of the general formula I or a pharmaceutically acceptable salt thereof for the preparation of a medicament for stimulating the release of growth hormone from the pituitary.

To those skilled in the art, it is well known that the current and potential uses of growth hormone in humans are varied and multitudinous. Thus, compounds of formula I can be administered for purposes stimulating release of growth hormone from the pituitary and would then have similar effects or uses as growth hormone itself. The uses of growth hormone may be summarized as follows: stimulation of growth hormone release in the elderly; prevention of catabolic side effects of glucocorticoids, prevention and treatment of osteoporosis, stimulation of the immune system, acceleration of wound healing, accelerating bone fracture repair, accelerating complicated fractures, e.g. distraction osteogenesis, treatment of wasting secondary to fractures, treatment of growth retardation, treating renal failure or insufficiency resulting from growth retardation, treatment of cardiomyopathy, treatment of chronic liver disease, treatment of thrombocytopenia, treatment of Crohn's disease, treatment of short bowel syndrome, treatment of chronic obstructive pulmonary disease (COPD), treatment of complications associated with transplantation, treatment of physiological short stature including growth hormone deficient children and short stature associated with chronic illness, treatment of obesity and growth retardation associated with obesity, treating growth retardation associated with the Prader-Willi syndrome and Turner's syndrome; accelerating the recovery and reducing hospitalization of burn patients; treatment of intrauterine growth retardation, skeletal dysplasia, hypercortisolism and Cushing's syndrome; induction of pulsatile growth hormone release; replacement of growth hormone in stressed patients, treatment of

- osteochondrodysplasias, Noonan's syndrome, schizophrenia, depressions, Alzheimer's disease, delayed wound healing and psychosocial deprivation, treatment of pulmonary dysfunction and ventilator dependency, attenuation of protein catabolic responses after major surgery, reducing cachexia and protein
- 5 loss due to chronic illness such as cancer or AIDS; treatment of hyperinsulinemia including nesidioblastosis, adjuvant treatment for ovulation induction; to stimulate thymic development and prevent the age-related decline of thymic function, treatment of immunosuppressed patients, improvement in muscle strength, mobility, maintenance of skin thickness, metabolic homeostasis, renal
- 10 homeostasis in the frail elderly, stimulation of osteoblasts, bone remodelling and cartilage growth, stimulation of the immune system in companion animals and treatment of disorder of aging in companion animals, growth promoter in livestock and stimulation of wool growth in sheep.
- 15 For the above indications the dosage will vary depending on the compound of formula I employed, on the mode of administration and on the therapy desired. However, generally dosage levels between 0.0001 and 100 mg/kg body weight daily are administered to patients and animals to obtain effective release of endogenous growth hormone. Usually, dosage forms suitable for oral, nasal,
- 20 pulmonal or transdermal administration comprise from about 0.0001 mg to about 100 mg, preferably from about 0.001 mg to about 50 mg of the compounds of formula I admixed with a pharmaceutically acceptable carrier or diluent.

25 Optionally, the pharmaceutical composition of the invention may comprise a compound of formula I combined with one or more compounds exhibiting a different activity, e.g., an antibiotic or other pharmacologically active material.

The route of administration may be any route which effectively transports the active compound to the appropriate or desired site of action, such as oral, nasal,

30 pulmonary, transdermal or parenteral, the oral route being preferred.

Apart from the pharmaceutical use of the compounds of formula I, they may be useful in vitro tools for investigating the regulation of growth hormone release.

Compounds of formula I may also be useful in vivo tools for evaluating the growth hormone releasing capability of the pituitary. For example, serum samples taken before and after administration of these compounds to humans can be assayed for growth hormone. Comparison of the growth hormone in each serum sample would directly determine the ability of the patients pituitary to release growth hormone.

Compounds of formula I may be administered to commercially important animals to increase their rate and extent of growth, and to increase milk production.

A further use of growth hormone secretagogue compounds of formula I is in combination with other secretagogues such as GHRP (2 or 6), GHRH and its analogues, growth hormone and its analogues or somatomedins including IGF-1 and IGF-2.

Pharmacological Methods

Compounds of formula I may be evaluated in vitro for their efficacy and potency to release growth hormone in rat pituitary primary cultures.

The isolation of rat pituitary cells is a modification of O. Sartor et al., Endocrinology **116**, 1985, pp. 952-957. Male albino Sprague-Dawley rats (250 +/- 25 grams) were purchased from Møllegaard, Lille Skensved, Denmark. The rats were housed in group cages (four animals/cage) and placed in rooms with 12 hour light cycle. The room temperature varied from 19-24°C and the humidity from 30 - 60%.

The rats were decapitated and the pituitaries dissected. The neurointermediate lobes were removed and the remaining tissue was immediately placed in icecold isolation buffer (Gey's medium (Gibco 041-04030) supplemented with 0.25% D-glucose, 2% non-essential amino acids (Gibco 043-01140) and 1% bovine serum albumine (BSA) (Sigma A-4503)). The tissue was cut into small pieces and transferred to isolation buffer supplemented with 3.8 mg/ml of trypsin (Worthington #3707 TRL-3) and 330 µg/ml of DNase (Sigma D-4527). This mixture was

incubated at 70 rotations/min for 35 min at 37°C in a 95/5% atmosphere of O₂/CO₂. The tissue was then washed three times in the above buffer. Using a standard pasteur pipette, the tissue was then aspirated into single cells. After dispersion, cells were filtered through a nylon filter (160 µm) to remove undigested
5 tissue. The cell suspension was washed 3 times with isolation buffer supplemented with trypsin inhibitor (0.75 mg/ml, Worthington #2829) and finally resuspended in culture medium; DMEM (Gibco 041-01965) supplemented with 25 mM HEPES (Sigma H-3375), 4 mM glutamine (Gibco 043-05030H), 0.075% sodium bicarbonate (Sigma S-8875), 0.1% non-essential amino acid, 2.5% fetal
10 calf serum (FCS, Gibco 011-06290), 3% horse serum (Gibco 034-06050), 10% fresh rat serum, 1 nM T₃ (Sigma T-2752) and 40 µg/l dexamethasone (Sigma D-4902) pH 7.3, to a density of 2 x 10⁵ cells/ml. The cells were seeded into microtiter plates (Nunc, Denmark), 200 µl/well, and cultured for 3 days at 37°C and 8% CO₂.

15 **Compound testing**

After culturing, the cells were washed twice with stimulation buffer (Hanks Balanced Salt Solution (Gibco 041-04020) supplemented with 1% BSA (Sigma A-4503), 0.25% D-glucose (Sigma G-5250) and 25 mM HEPES (Sigma H-3375) pH
20 7.3) and preincubated for 1 hour at 37°C. The buffer was exchanged with 90 µl stimulation buffer (37°C). Ten µl test compound solution was added and the plates were incubated for 15 min at 37°C and 5% CO₂. The medium was decanted and analyzed for GH content in an rGH SPA test system.

25 All compounds were tested in doses ranging from 10 pM to 100 µM. A dose-response relation was constructed using the Hill equation (Fig P, Biosoft). The efficacy (maximal GH released, E_{max}) was expressed in % of the E_{max} of GHRP-6. The potency (EC₅₀) was determined as the concentration inducing half maximal stimulation of the GH release.

30

Compounds of formula I may be evaluated for their metabolic stability.

Compounds were dissolved at a concentration of 1 µg/µl in water. 25 µl of this solution is added to 175 µl of the respective enzyme-solution (resulting in an

enzyme:substrate ratio (w/w) of approximately 1:5). The solution is left at 37°C overnight. 10 µl of the various degradation solutions is analyzed against a corresponding zero-sample using flow injection electrospray mass spectrometry (ESMS) with selected ion monitoring of the molecular ion. If the signal has
5 decreased more than 20% compared to the zero-sample, the remainder of the solution is analyzed by HPLC and mass spectrometry in order to identify the extent and site(s) of degradation precisely.

Several standard peptides (ACTH 4-10, Angiotensin 1-14 and Glucagon) have
10 been included in the stability tests in order to verify the ability of the various solutions to degrade peptides.

Standard peptides (angiotensin 1-14, ACTH 4-10 and glucagon) were purchased from Sigma, MO, USA)
15

Enzymes (trypsin, chymotrypsin, elastase aminopeptidase M and carboxypeptidase Y and B) were all purchased from Boehringer Mannheim GmbH (Mannheim, Germany)

20 Pancreatic enzyme mix: trypsin, chymotrypsin and elastase in 100 mM ammoniumbicarbonate pH 8.0 (all concentrations 0.025 µg/µl).

Carboxypeptidase mix: carboxypeptidase Y and B in 50 mM ammoniumacetate pH 4.5 (all concentrations 0.025 µg/µl).
25

Aminopeptidase M solution: aminopeptidase M (0.025 µg/µl) in 100 mM ammoniumbicarbonate pH 8.0

Mass spectrometric analysis was performed using two different mass
30 spectrometers. A Sciex API III triple quadrupole LC-MS instrument (Sciex instruments, Thornhill, Ontario) equipped with an electrospray ion-source and a Bio-Ion 20 time-of-flight Plasma Desorption instrument (Bio-Ion Nordic AB, Uppsala, Sweden).

Quantification of the compounds (before and after degradation) was done on the API III instrument using single ion monitoring of the molecular ion in question with flow injection of the analyte. The liquid flow (MeOH:water 1:1) of 100 μ l/min was controlled by an ABI 140B HPLC unit (Perkin-Elmer Applied Biosystems Divisions, Foster City, CA). The instrument parameters were set to standard operation conditions, and SIM monitoring was performed using the most intense molecular ion (in most cases this corresponded to the doubly charged molecular ion).

Identification of degradation products furthermore involved the use of plasma desorption mass spectrometry (PDMS) with sample application on nitrocellulose coated targets and standard instrumental settings. The accuracy of the hereby determined masses is generally better than 0.1%.

Separation and isolation of degradation products was done using a HY-TACH C-18 reverse phase 4.6x105 mm HPLC column (Hewlett-Packard Company, Palo Alto, CA) with a standard acetonitril: TFA separation gradient. The HPLC system used was HP1090M (Hewlett-Packard Company, Palo Alto, CA).

Peptide derivative	MW/SIM ion (amu)	Carboxy-peptidase mix	Pan. enzyme mix
Standards			
ACTH 4-10	1124.5/562.8	+	-
Glucagon	3483/871.8	-	-
Insulin (B23-29)	859.1/430.6		
Angiotensin 1-14	1760.1/881.0	-	-
GHRP-2	817.4/409.6	-	-
GHRP-6	872.6/437.4	-	-

20 +: Stable (less than 20% decrease in SIM signal after 24 h in degradation solution)
 -: Unstable (more than 20% decrease in SIM signal after 24 h in degradation solution)

Any novel feature or combination of features described herein is considered essential to this invention.

Chemical Methods


5

Some compounds of formula I or intermediates may be synthesized by one of the reactions sequences shown. Other examples may require different synthetic approaches, depending of the type of compound. The specific synthesis of each example is therefore given in the description of each example.

10

Method A

Scheme 1




- 5 An aldehyde 1 may be reacted with an ester 2 under reductive conditions such as e.g. sodium cyano borohydride to give an ester 3. After deprotection of the amino group a ring closure may take place to give a piperazinone 4. A reaction with a suitable protected amino acid 5 with a coupling reagent known in the art such as e.g. 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride or a
- 10 combination of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride

and 1-hydroxybenzotriazole or 1-hydroxy-7-azabenzotriazole in an appropriate solvent such as e.g. N,N-dimethylformamide or dichloromethane may produce the piperazinone **6**. After deprotection carried out with a method known in the art and described by e.g. T. W. Greene (Protective Groups in Organic Synthesis, 2. ed., John Wiley and Sons, New York 1991) e.g. trifluoroacetic acid or hydrogen chloride in ethyl acetate, the resulting amine **7** may be coupled with a suitable protected amino acid **8** with a coupling reagent known in the art such as e.g. 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride or a combination of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and 1-hydroxybenzotriazole or 1-hydroxy-7-azabenzotriazole in an appropriate solvent such as e.g. N,N-dimethylformamide or dichloromethane and successive deprotection with a method known in the art and described by e.g. T. W. Greene (Protective Groups in Organic Synthesis, 2. ed., John Wiley and Sons, New York 1991) e.g. trifluoroacetic acid or hydrogen chloride in ethyl acetate to give **9**, which is a compound of formula I.

Method B

Scheme 2



- 5 12 13
- A diazepinone may be synthesized as described in scheme 2 and incorporated into the peptide in a similar fashion as in scheme 1 by reductive alkylation of a carbonyl-compound 11 onto an amino acid ester 10 under dehydrating conditions such as e.g. molecular sieves and sodium cyanoborohydride in a solvent such as e.g. methanol and subsequent removal of a protecting group from 12. This may result in cyclization of the intermediate to form 13. 13 may be used as an intermediate, which may be reacted in the same manner as described in scheme 1 for the intermediate 4 to give a compound of formula I.
- 10

EXAMPLES:

The process for preparing compounds of formula I and preparations containing them is further illustrated in the following examples, which however, are not to be construed as limiting.

The structures of the compounds are confirmed by either elemental analysis (MA) nuclear magnetic resonance (NMR) or mass spectrometry (MS). NMR shifts (δ) are given in parts per million (ppm) and only selected peaks are given. mp is melting point and is given in °C. Column chromatography was carried out using the technique described by W.C. Still et al, J. Org. Chem. 1978, 43, 2923-2925 on Merck silica gel 60 (Art 9385). Compounds used as starting materials are either known compounds or compounds which can readily be prepared by methods known per se.

15

Abbreviations:

TLC: thin layer chromatography
DMSO: dimethylsulfoxide
min: minutes
20 h: hours

HPLC-Analysis:**Method A1.**


25

The RP-analysis was performed using UV detections at 214, 254, 276, and 301 nm on a 218TP54 4.6 mm x 250 mm 5 μ C-18 silica column (The Separations Group, Hesperia), which was eluted at 1 mL/min at 42°C. The column was equilibrated with 5% acetonitrile in a buffer consisting of 0.1 M ammonium sulfate, which was adjusted to pH 2.5 with 4M sulfuric acid. after injection the sample was eluted by a gradient of 5% to 60% acetonitrile in the same buffer during 50 min.

30

Example 1


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxo-piperazin-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.



5

(2R)-2-(2-(tert-Butoxycarbonylamino)ethylamino)-3-phenylpropionic acid methyl ester.

10




- 15 (2R)-2-Amino-3-phenylpropionic acid methyl ester (3.2 g; 15.0 mmol) and (2-oxoethyl)carbamic acid tert-butyl ester (3.2 g; 20.0 mmol) (prepared as in Dueholm et al, Org. Prep. Proced. Int. 25(4), 457-61 (1993)) were dissolved in a mixture of methanol (100 ml) and acetic acid (5 ml). Molecular sieves (3 Å; 100 g) were added and the reaction mixture was cooled to 0°C. Sodium cyanoborohydride (1.38 g; 22.0 mmol) was added portionwise and the reaction mixture was left without stirring for 12 h. The reaction mixture was filtered and water (100 ml), aqueous sodium hydrogencarbonate/sodium carbonate (10%; 100 ml; pH 9) and methylene chloride (50 ml) were added to the filtrate. The aqueous phase was extracted with methylene chloride (2 x 50 ml) and the combined organic phases were dried over magnesium sulfate. The organic
- 20

phase was evaporated in vacuo and the residue was chromatographed on silica (4 x 40 cm) using ethyl acetate/heptane (3:2) as eluent to afford 2.07 g of (2R)-2-(2-(tert-butoxycarbonylamino)ethylamino)-3-phenylpropionic acid methyl ester.

- 5 ¹H-NMR (CDCl₃): δ 1.44 (s, 9H); 2.55 (m, 1H); 2.72 (m, 1H); 2.90 (dd, 1H); 2.97 (dd, 1H); 3.11 (m, 2H); 3.47 (t, 1H); 3.68 (s, 3H); 7.15-7.32 (arom., 5H);


- 10 (2R)-2-(2-Aminoethylamino)-3-phenylpropionic acid methyl ester.



- (2R)-2-(2-(tert-Butoxycarbonylamino)ethylamino)-3-phenylpropionic acid methyl ester (1.32 g; 4.094 mmol) was dissolved in methylene chloride (10 ml) and trifluoroacetic acid (10 ml) was added. The reaction mixture was stirred for 1 hour at room temperature. The solvent was evaporated in vacuo and the residue was dissolved in methylene chloride (5 ml) and evaporated in vacuo. Methylene chloride (5 ml) was added and evaporated in vacuo to afford 1.14 g of (2R)-2-(2-aminoethylamino)-3-phenylpropionic acid methyl ester as a trifluoroacetate salt.
- 15
- 20

¹H-NMR (DMSO-d₆): δ 3.03-3.39 (m, 6H); 3.66 (s, 3H); 4.45 (dd, 1H); 7.21-7.39 (arom., 5H)


- 25 (3R)-3-Benzylpiperazin-2-one.



(2R)-2-(2-Aminoethylamino)-3-phenylpropionic acid methyl ester (0.67 g; 1.49 mmol) and triethylamine (1 ml) were dissolved in methylene chloride (10 ml) and stirred for 48 hours at room temperature. The reaction mixture was evaporated and the residue was chromatographed on silica (2.5 x 30 cm) using a 7% solution of ammonia in ethanol/methylene chloride (1:9) as eluent to afford 0.18 g of (3R)-3-benzylpiperazin-2-one.

¹H-NMR (CDCl₃): δ 1.72 (s(br), 1H); 2.81-3.48 (m, 6H); 3.62 (dd, 1H); 6.92 (s(br), 1H); 7.19-7.36 (arom., 5H).

N-((1R)-2-((2R)-2-Benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylcarbamic acid tert-butyl ester.



15


(2R)-2-(tert-Butoxycarbonylmethylamino)-3-(2-naphthyl)propionic acid (prepared as in J.R. Coggins, N.L. Benoit, Can.J.Chem 49, 1968 (1971)) (0.327 g; 0.995 mmol) was dissolved in methylene chloride (10 ml). 1-Hydroxy-7-azabenzotriazole (0.135 g; 0.995 mmol) and N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (0.199 g; 1.042 mmol) were added and the reaction mixture was stirred for 10 min at room temperature. (3R)-3-Benzylpiperazin-2-one (0.180 g;

20

0.947 mmol) and diisopropylethylamine (0.134 g; 1.042 mmol) were added and the reaction mixture was stirred for 12 hours at room temperature. The reaction mixture was washed with 10% sodium hydrogensulfate (20 ml), saturated sodium hydrogencarbonate (20 ml), dried over magnesium sulfate and evaporated in vacuo. The residue was chromatographed on silica (2.5 x 25 cm) using ethyl acetate as eluent to afford 0.405 g of N-((1R)-2-((2R)-2-benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylcarbamic acid tert-butyl ester.

¹H-NMR (CDCl₃): δ 1.12, 1.25, 1.38 (three s, 9H); 2.54, 2.55, 2.78 (three s, 3H); 4.07, 4.42, 4.65, 4.79, 5.01, 5.30 (six m; 2H); 7.10-7.78 (arom., 12 H) (mixture of rotamers, selected peaks).

(3R)-3-Benzyl-4-((2R)-2-methylamino-3-(2-naphthyl)propionyl)piperazin-2-one.




15

N-((1R)-2-((2R)-2-Benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylcarbamic acid tert-butyl ester (0.405 g; 0.808 mmol) was dissolved in methylene chloride (5 ml) and trifluoroacetic acid (5 ml) and stirred 10 min at room temperature. Water (10 ml) and sodium hydrogencarbonate were added to pH 8. The reaction mixture was extracted with methylene chloride (2 x 30 ml), dried over magnesium sulfate and evaporated in vacuo to afford 0.302 g of (3R)-3-benzyl-4-((2R)-2-methylamino-3-(2-naphthyl)propionyl)piperazin-2-one.


¹H-NMR (CDCl₃): δ 2.19, 2.28 (two s, 3H); 3.72, 4.66 (two dd, 1H); 4.96, 5.32 (two dd, 1H) (mixture of rotamers, selected peaks).

3-Hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester.



- At 0°C, ethyl chloroformate (1.10 ml, 11.5 mmol) was given dropwise to a solution of 3-tert-butoxycarbonylamino-3-methylbutanoic acid (2.50 g, 11.5 mmol) (cf. Schoen et al. J. Med. Chem., 1994, vol. 37, p. 897-906, and references cited herein) and triethylamine (1.92 ml, 13.8 mmol) in tetrahydrofuran (10 ml). The solution was stirred for 40 min at 0°C. The formed precipitate was filtered off and washed with tetrahydrofuran (20 ml). The liquid was immediately cooled to 0°C. A 2M solution of lithium borohydride in tetrahydrofuran (14.4 ml, 28.8 mmol) was added dropwise. The solution was stirred at 0°C for 2 h, and then warmed to room temperature over a period of 4 h. It was cooled to 0°C and methanol (5 ml) was added. 1N Hydrochloric acid (100 ml) was added and the solution was extracted with ethyl acetate (2 x 100 ml, 3 x 50 ml). The combined organic extracts were washed with saturated sodium hydrogencarbonate solution (100 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was chromatographed on silica (110 g) with ethyl acetate/heptane (1:2) to give 1.84 g of 3-hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester.
- ¹H-NMR (CDCl₃): δ 1.33 (s, 6 H); 1.44 (s, 9 H); 1.88 (t, 2 H); 1.94 (br, 1 H); 3.75 (q, 2 H); 4.98 (br, 1 H).

3-(tert-Butoxycarbonylaminomethyl)-3-methylbutanal.




- At -78°C dimethylsulfoxide (1.22 ml, 17.2 mmol) was added to a solution of oxalyl chloride (1.1 ml, 12.9 mmol) in dichloromethane (15 ml). The mixture was stirred for 15 min at -78°C. A solution of 3-hydroxy-1,1-dimethylpropylcarbamic acid tert-butyl ester (1.75 g, 8.6 mmol) in dichloromethane (10 ml) was added dropwise over a period of 15 min. The solution was stirred at -78°C for another 15 min. Triethylamine (6.0 ml, 43 mmol) was added. The solution was stirred at -78°C for

5 min and then warmed to room temperature. The solution was diluted with dichloromethane (100 ml) and extracted with 1N hydrochloric acid (100 ml). The aqueous phase was extracted with dichloromethane (50 ml). The combined organic layers were washed with saturated sodium hydrogencarbonate solution (100 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by column chromatography on silica (140 g) with ethyl acetate/heptane (1:3) to give 1.10 g of 3-(tert-butoxycarbonylaminoethyl)-3-methylbutanal.

¹H-NMR (CDCl₃): δ 1.39 (s, 6 H); 1.45 (s, 9 H); 2.85 (d, 2 H); 4.73 (br. 1 H); 9.80 (t, 1 H).


Ethyl (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoate.



Triethylphosphonoacetate (1.96 ml, 9.8 mmol) was dissolved in tetrahydrofuran (30 ml). Potassium tert-butoxide (1.10 g, 9.8 mmol) was added. The solution was stirred for 40 min. at room temperature. A solution of 3-(tert-butoxycarbonylaminoethyl)-3-methylbutanal (1.10 g, 5.5 mmol) in tetrahydrofuran (6 ml) was added slowly. The solution was stirred at room temperature for 75 min. It was diluted with ethyl acetate (100 ml) and 1N hydrochloric acid (100 ml). The phases were separated. The aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic layers were washed with saturated sodium hydrogencarbonate solution (60 ml) and dried over magnesium sulfate. The solvent was removed in vacuo. The crude product was purified by column chromatography on silica (90 g) with ethyl acetate/heptane (1:4) to give 1.27 g of ethyl (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoate.

¹H-NMR (CDCl₃): δ 1.30 (s, 6 H); 1.30 (t, 3 H); 1.46 (s, 9 H); 2.62 (d, 2 H); 4.27 (q, 2 H); 4.42 (br, 1 H); 5.88 (d, 1 H); 6.94 (td, 1 H).


(2E)-5-(tert-Butyloxycarbonylamino)-5-methylhex-2-enoic acid.



Ethyl (2E)-5-(tert-butoxycarbonylamino)-5-methylhex-2-enoate (1.23 g, 4.54 mmol) was dissolved in dioxane (20 ml). Lithium hydroxide (0.12 g, 5.00 mmol) was added as a solid. Water (10 ml) was added, until a clear solution was obtained. The solution was stirred for 16 h at room temperature. The solution was diluted with water (70 ml) and was extracted with tert-butyl methyl ether (2 x 100 ml). The aqueous phase was acidified with 1N sodium hydrogensulfate solution (pH 1) and extracted with tert-butyl methyl ether (3 x 70 ml). These organic layers were combined and dried over magnesium sulfate. The solvent was removed in vacuo to give 1.05 g of (2E)-5-(tert-butyloxycarbonylamino)-5-methylhex-2-enoic acid. The crude product was used for further syntheses.

¹H-NMR (DMSO-d₆): δ 1.15 (s, 6 H); 1.35 (s, 9 H); 2.53 (d, 2 H); 5.75 (d, 1 H); 6.57 (br, 1 H); 6.75 (td, 1 H); 12.15 (s, 1 H).

((3E)-4-(N-((1R)-2-((2R)-2-Benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylcarbamoyl)-1,1-dimethylbut-3-enyl)carbamic acid tert-butyl ester.



(2E)-5-(tert-Butyloxycarbonylamino)-5-methylhex-2-enoic acid (0.145 g; 0.599 mmol) was dissolved in methylene chloride. 1-Hydroxy-7-azabenzotriazole (0.081 g; 0.599 mmol) and N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (0.126 g; 0.658 mmol) were added and the reaction mixture was stirred for 15 min. (3R)-3-Benzyl-4-((2R)-2-methylamino-3-(2-naphthyl)propionyl-

piperazin-2-one (0.24 g; 0.599 mmol) and diisopropylethylamine (0.077 g; 0.599 mmol) were added and the reaction mixture was stirred for 12 hours at room temperature. The reaction mixture was washed with sodium hydrogensulfate (100 ml) and sodium hydrogencarbonate (100 ml). The organic phase was evaporated in vacuo and the residue was chromatographed on silica (2.5 x 20 cm) using ethyl acetate as eluent to afford 0.332 g of ((3E)-4-(N-((1R)-2-((2R)-2-benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)-methyl)-2-oxoethyl)-N-methylcarbamoyl)-1,1-dimethylbut-3-enyl)carbamic acid tert-butyl ester.

¹H-NMR (CDCl₃):) (mixture of rotamers, selected peaks for major rotamer) δ 1.24 (s, 3H); 1.27 (s, 3H); 1.42 (s, 9H); 2.72 (s, 3H); 5.30 (dd, 1H); 5.81 (dd, 1H); 6.18 (d, 1H); 6.82 (m, 1H).

((3E)-4-(N-((2R)-2-((2R)-2-benzyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxo-ethyl)-N-methylcarbamoyl)-1,1-dimethylbut-3-enyl)carbamic acid tert-butyl ester (0.32 g; 0.511 mmol) was dissolved in methylene chloride (4 ml) and trifluoroacetic acid (4 ml) was added. The reaction mixture was stirred 12 min at room temperature. Methylene chloride (50 ml) and water (50 ml) were added. Sodium hydrogencarbonate was added until pH 8. The organic phase was dried over magnesium sulfate and evaporated in vacuo to afford 0.232 g of the title compound.


¹H-NMR (CDCl₃): δ 1.14, 1.15, 1.22, 1.23 (four s, 6H); 4.05, 4.60, 4.76, 5.12, 5.38, 5.81 (six dd, 2H); 6.18, 6.22 (d, 1H); 6.70, 6.81 (two m, 1H) (mixture of rotamers, selected peaks).

ESMS: m/z 527 (M+H)⁺

HPLC (A1): *t*_r: 31.03 min.


Example 2

(2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.



5

(1,1-Dimethyl-3-oxobutyl)carbamic acid tert-butyl ester:



10

Diacetonamine hydrogen oxalate (30.0 g; 146 mmol) was suspended in tetrahydrofuran (400 ml). An aqueous solution of sodium hydroxide (1 N; 146 ml) was added. Di-tert-butyl dicarbonate (38.3 g; 175 mmol) was dissolved in tetrahydrofuran (100 ml) and added dropwise to the reaction mixture. The reaction mixture was stirred for 2 hours at room temperature. Sodium hydroxide (1 N; 146 ml) was added and the reaction mixture was stirred for 12 h at room temperature. Water (200 ml) and ethyl acetate (200 ml) were added. The aqueous phase was extracted with ethyl acetate (4 x 200 ml). The combined organic phases were dried over magnesium sulfate and the solvent was removed in vacuo. The residue was chromatographed on silica (6 x 40 cm) using ethyl acetate/heptane (1:3) as eluent to afford 28.4 g of (1,1-dimethyl-3-oxobutyl)carbamic acid tert-butyl ester.


15

20

¹H-NMR (CDCl₃) δ 1.34 (s, 6H); 1.42 (s, 9H); 2.14 (s, 3H); 2.86 (s, 2H); 4.85 (s, 1H).

25

(E)-5-tert-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid ethyl ester:




Triethyl phosphonoacetate (4.7 g; 20.9 mmol) was dissolved in tetrahydrofuran
 5 (36 ml). Potassium tert-butoxide (2.3 g; 20.9 mmol) was added and the reaction
 mixture was stirred for 40 min at room temperature. (1,1-Dimethyl-3-
 oxobutyl)carbamic acid tert-butyl ester (2.5 g; 11.6 mmol) was dissolved in
 tetrahydrofuran (15 ml) and added dropwise to the reaction mixture which was
 heated to reflux for 12 h. Ethyl acetate (100 ml) and hydrochloric acid (1 N; 100
 10 ml) were added and the phases were separated. The aqueous phase was
 extracted with ethyl acetate (3 x 50 ml). The combined organic phases were
 washed with an aqueous solution of sodium hydrogen carbonate (saturated; 100
 ml), dried over magnesium sulfate and evaporated in vacuo. The residue was
 chromatographed on silica (3 x 40 cm) using ethyl acetate/heptane (1:2) as eluent
 15 to afford 2.0 g of (E)-5-tert-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid
 ethyl ester.

¹H-NMR (CDCl₃): δ 1.25 (t, 3H); 1.30 (s, 6H); 1.44 (s, 9H); 2.21 (s, 3H); 2.58 (s,
 2H); 4.14 (q, 2H); 4.48 (s, 1H); 5.65 (s, 1H).

20

(2E)-5-tert-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid:



(E)-5-tert-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid ethyl ester (1.95 g;
 6.83 mmol) was dissolved in 1,4-dioxane (25 ml) and water (15 ml). Lithium
 hydroxide (0.18 g; 7.52 mmol) was added and the reaction mixture was stirred for
 12 h at room temperature. Water (150 ml) and tert-butyl methyl ether (150 ml) was
 added. The aqueous phase was diluted with an aqueous solution of sodium
 25 hydrogensulfate (10 %) until pH 2.5 and extracted with tert-butyl methyl ether (3 x
 30 100 ml). The combined organic phases were dried over magnesium sulfate and

evaporated in vacuo. The residue was recrystallized from heptane (20 ml) to afford 0.6 g of (2E)-5-tert-butoxycarbonylamino-3,5-dimethylhex-2-enoic acid.

¹H-NMR (CDCl₃) δ 1.29 (s, 6H); 1.44 (s, 9H); 2.23 (s, 3H); 2.62 (s, 2H); 4.45 (s,
5 1H); 5.66 (s, 1H).

The remaining synthesis proceeded similarly as in Example 1 to afford the title compound. (2E)-5-tert-Butoxycarbonylamino-3,5-dimethylhex-2-enoic acid was used in the last coupling step instead of (2E)-5-tert-butoxycarbonylamino-5-methylhex-2-enoic acid.
10

¹H-NMR (CDCl₃) (selected peaks for major rotamer) δ 1.37 (s, 6H); 2.09 (s, 2H); 2.50 (s, 3H); 2.78 (s, 3H); 4.67 (dd, 1H); 5.05 (dd, 1H); 5.89 (s, 1H).


ESMS: m/z 541.7 (M+H)⁺

15 HPLC (A1): r_t: 32.40 min


Example 3

(2E)-5-Methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxo-piperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.

5



(2E)-5-(N-(tert Butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid.



10

(2E)-5-(tert-Butyloxycarbonylamino)-5-methylhex-2-enoic acid (5.00 g ; 20.6 mmol) was dissolved in tetrahydrofuran (70 ml). Methyl iodide (10.3 ml; 164 mmol) was added and the solution was cooled to 0° C. Sodium hydride (60% in oil) (2.07 g; 61.6 mmol) was added in portions and the solution was stirred at room temperature for four days. Ethyl acetate (70 ml) and water (60 ml) were added dropwise and the solvent was removed in vacuo. The crude product was dissolved in water (40 ml) and diethyl ether (40 ml). The organic phase was washed with a saturated aqueous solution of sodium hydrogencarbonate (30 ml). The aqueous phases were mixed and 5% aqueous citric acid was added to pH 3. The aqueous phase was extracted with ethyl acetate (4 x 50 ml). The combined organic extracts were washed with water (2 x 40 ml), an aqueous solution of sodium thiosulfate (5%; 40 ml), water (40 ml), dried over MgSO₄ and the solvent was removed in vacuo. The residue was dissolved in ethyl acetate (45 ml) and washed with an aqueous solution of sodium hydrogensulfate (10%;

3 x 30 ml), dried over magnesium sulfate and concentrated in vacuo to give 4.00 g of (2E)-5-(N-(tert-butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid.


¹H-NMR (CDCl₃) δ 1.38 (s, 6H), 1.45 (s, 9H); 2.80 (d, 2H); 2.85 (s, 3H); 5.88 (d, 1H); 7.01 (q, 1H).

The remaining synthesis proceeded similarly as in Example 1 substituting (2E)-5-(tert-butyloxycarbonylamino)-5-methylhex-2-enoic acid with (2E)-5-(N-(tert-butoxycarbonyl)-N-methylamino)-5-methylhex-2-enoic acid in the last coupling step to afford the title compound.

¹H-NMR (CDCl₃) (selected peaks for major rotamer) δ 1.38 (s, 3H); 1.44 (s, 3H); 2.69 (s, 3H); 2.88 (s, 3H); 4.44 (dd, 1H); 4.52 (dd, 1H); 6.34 (d, 1H).
ESMS: m/z 541.9 (M+H)⁺
HPLC (A1): t_r: 31.37 min

Example 4

(2E)-5-Methyl-5-methylaminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.



This compound was prepared using the procedure in Example 1 and 3. L-Boc-alaninal was used instead of Boc-glycinal.

¹H-NMR (CDCl₃): (selected peaks for major rotamer) δ 1.21 (d, 3H); 1.32 (s, 3H); 1.40 (s, 3H); 2.63 (s, 3H); 2.88 (s, 3H); 4.42 (dd, 1H); 4.60 (dd, 1H); 6.26 (d, 1H).


ESMS: m/z 555.9 (M+H)⁺

HPLC (A1): t_r : 32.30 min.


Example 5

5

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide



10



15 (3R)-4-tert-Butyloxycarbonyl-3-benzylpiperazin-2-one


(3R)-3-benzylpiperazin-2-one (from Example 1) (1.3 g) was dissolved in THF (20 ml) and aqueous sodium hydroxide (1 M, 7 ml) was added. Boc-anhydride (1.8 g) was dissolved in THF (10 ml) and added dropwise. The mixture was stirred overnight. THF was removed in vacuo. Water (10 ml) was added and the aqueous phase was extracted with ethyl acetate (2 x 50 ml). The combined organic phases were dried (magnesium sulphate) and the solvent was removed in vacuo and the residue was chromatographed on Silica using ethyl acetate as eluent to afford 750 mg of 3R-4-tert-butyloxycarbonyl-3-benzylpiperazin-2-one.

20

¹H-NMR (CDCl₃) (major rotamer): δ 1.31 (s, 9H); 2.85 (t, 1H); 3.10 (m, 2H); 3.25 (m, 2H); 3.42 (t, 2H); 4.80 (m, 1H); 6.40 (s, 1H); 7.12-7.35 (5 arom.H).

5

(3R)-4-tert-Butyloxycarbonyl-1-methyl-3-benzylpiperazin-2-one.



10 (3R)-4-tert-Butyloxycarbonyl-3-benzylpiperazin-2-one (see Example 1) (400 mg; 1.4 mmol) was added to a mixture of KOH (300 mg; 5.5 mmol) and DMSO (3 ml). Methyl iodide (400 mg 2.7 mmol) was added and the mixture was stirred for 1h. Methylene chloride (20 ml) was added and the organic phase was washed with water (5 x 10 ml). The solvent was removed in vacuo to afford 220 mg of (3R)-4-tert-butyloxycarbonyl-1-methyl-3-benzylpiperazin-2-one.

15

¹H-NMR (CDCl₃) δ 1.21 (s, 9H); 2.97 (s, 3H); 4.13 (dd, 1H).

The remaining steps were performed similarly as in example 1 to afford the title compound.

20

¹H-NMR (CDCl₃) (selected peaks for major rotamer) δ 1.15 (s, 6H); 2.30 (d, 2H); 2.54 (s, 3H); 2.74 (s, 3H); 4.05 (dd, 1H); 5.72 (dd, 1H); 6.19 (d, 1H).

ESMS: m/z 541.9 (M+H)⁺


HPLC (A1): t_r: 27.70 min

25

Example 6

(2E)-5-Methyl-5-aminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide.

5



The title compound was prepared using the procedure in Example 1 and 4.


10 ¹H-NMR (CDCl₃): δ 1.09 (d, 3H); 1.31 (s, 6H); 2.74 (s, 3H); 5.12 (dd, 1H); 6.18 (d, 1H).

ESMS: m/z 541.7 (M+H)⁺

HPLC (A1): t_r: 32.12 min.

15 Example 7

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-methyl-N-((1R)-2-(2-naphthyl)-1-((2R)-3-oxo-2-((2-thienyl)methyl)piperazine-1-carbonyl)ethyl)amide



20


The title compound was prepared using the procedure in Schoen et al. J. Med. Chem., 1994, vol. 37, p. 897-906 substituting isobutylene with methylene cyclobutane and D-phenylalanine with D-thienylalanine.

- 5 ¹H-NMR (CDCl₃): δ 3.01(s, 3H); 4.76 (dd, 1H); 5.15 (t, 1H); 6.34 (d, 1H)
 ESMS: m/z 546.0 (M+H)⁺
 HPLC (A1): r_t: 31.05 min.

Example 8

10

(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-3-oxo-piperazin-1-yl)-1-((biphenyl-4-yl)methyl)-2-oxoethyl)-N-methylamide.



15


The title compound was prepared using the procedure in Example 1 using 4,4'-biphenylalanine instead of 2-naphthylalanine.

- 20 ¹H-NMR (CDCl₃): δ 1.18 (s, 6H); 2.77 (s, 1H); 5.31 (dd, 1H); 5.72 (dd, 1H); 6.21 (dd, 1H).
 ESMS: m/z 554.1 (M+H)⁺
 HPLC (A1): r_t: 34.62

Example 9

25

(2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-2-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide




- The title compound was prepared as in Schoen et al. J. Med. Chem., 1994, vol. 37, p. 897-906 substituting isobutylene with methylene cyclobutane and Boc-D-alaninal instead of Boc-glycinal.

- ¹H-NMR (CDCl₃): δ (selected peaks) 1.11 (d, 3H); 2.66 (s, 3H); 4.97 (dd, 1H); 5.85 (dd, 1H); 6.24 (d, 1H); 6.86 (m, 1H)
ESMS: m/z 554.0 (M+H)⁺
HPLC (A1): t_r: 32.3 min.

Example 10

- (2E)-5-Methyl-5-methylamino-hex-2-enoic acid N-methyl-N-((1R)-1-(2-naphthyl)methyl-2-oxo-2-((2R)-3-oxo-2-((2-thienyl)methyl)piperazin-1-yl)ethyl)amide



- The title compound was prepared similarly as in Example 1 substituting D-phenylalanine with D-thienylalanine.

¹H-NMR (CDCl₃): δ 1.38 (s, 3H); 1.41 (s, 3H); 2.65 (s, 3H); 2.91 (s, 3H); 4.47 (dd, 1H);

ESMS: m/z 547.0 (M+H)⁺


HPLC (A1): t_r: 31.15 min.

5

Example 11


(2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxoethyl)-N-methylamide.

10



(3R)-3,4-Dibenzylpiperazin-2-one.

15




20

(3R)-3-Benzylpiperazin-2-one (4.1 g; 21.6 mmol, prepared as described in Example 1) was dissolved in methanol (40 ml). Benzaldehyde (2.3 g; 21.6 mmol) was added followed by 4 Å molsieves (2 g) and the mixture was stirred under an atmosphere of nitrogen for ½ h. Acetic acid (1.2 ml) was added and sodium cyanoborohydride (2.0 g; 32.4 mmol) was added portion wise during 20 minutes. The reaction mixture was stirred overnight and the filtered. From the filtrate the solvent was evaporated and the residue was dissolved into ethyl acetate (15 ml) and purified on silica gel using ethyl acetate as eluent. This afforded 2.6 g of (3R)-3,4-dibenzylpiperazin-2-one.

25

¹H-NMR (CDCl₃): δ 2.50 (m, 1H); 2.95 (m, 1H); 3.08 (m, 1H); 3.15-3.45 (m, 4H); 3.48 (d, 1H); 3.96 (d, 1H); 6.18 (s, 1H); 7.1-7.3 (m, 10H).


(2R)-1,2-Dibenzylpiperazine.



5 A solution of (3R)-3,4-dibenzylpiperazin-2-one (2.5 g; 9.0 mmol) in dry tetrahydrofuran (30 ml) was placed under an atmosphere of nitrogen and stirred on an ice-bath. Lithium aluminiumhydride (5.0 ml; 1.0 M in tetrahydrofuran) was added dropwise during 15 minutes. The mixture was then heated at reflux temperature for 2 h and then allowed to cool. Another portion of lithium aluminiumhydride (4.0 ml; 1.0 M in tetrahydrofuran) was added and the mixture
10 was refluxed for 1 h. The mixture was allowed to cool and then quenched with water and a 4 N sodium hydroxide solution. Ethyl acetate (50 ml) was added and the mixture was filtered. The solvent was evaporated in vacuo to give 2.0 g of (2R)-1,2-dibenzylpiperazine.

15 ¹H-NMR (CDCl₃): δ 2.20 (m, 1H); 2.55-2.82 (m, 7H); 3.15 (dd, 1H); 3.42 (d, 1H); 4.10 (d, 1H); 7.1-7.4 (m, 10H).


(2R)-1,2-Dibenzyl-4-methanesulfonylpiperazine.



20 To a stirred solution of (2R)-1,2-dibenzylpiperazine (2.0 g; 7.5 mmol) in dichloromethane (30 ml), methanesulfonyl chloride (0.86 g; 7.5 mmol) was added and the mixture was stirred at ambient temperature for 1 h. Potassium carbonate (1.1 g) was added and stirring was continued for another hour. Dichloromethane (20 ml) and water (10 ml) were added and the phases were
25 separated. The organic phase was washed with water (10 ml) and dried over magnesium sulfate. The solvent was evaporated in vacuo to give 2.2 g of (2R)-1,2-dibenzyl-4-methanesulfonylpiperazine.

¹H-NMR (CDCl₃): δ 2.58 (m, 1H); 2.72 (s, 3H); 2.85-3.10 (m, 6H); 3.27 (m, 1H); 3.45 (m, 1H); 3.72 (d, 1H); 3.95 (m, 1H); 7.1-7.4 (m, 10H).


(3R)-3-Benzyl-1-methanesulfonylpiperazine hydrochloride.



(2R)-1,2-Dibenzyl-4-methanesulfonylpiperazine (2.2 g; 6.4 mmol) was dissolved in dry 1,2-dichloroethane (20 ml) under an atmosphere of nitrogen. The mixture was placed on an ice-bath and a solution of α-chloroethyl chloroformate (1.0 g; 7.04 mmol) in dry 1,2-dichloroethane (10 ml) was added keeping the temperature between 0 and 5°C. When addition is complete stirring is continued for 15 minutes on the ice-bath. The reaction mixture is heated slowly to 90°C on an oil-bath and heated at reflux temperature for 1 h. The major part of the solvent was allowed to evaporate and the residue was allowed to cool on an ice-bath. Methanol (20 ml) was added and the mixture was heated at reflux temperature for 1 h. The mixture was allowed to cool to ambient temperature, diethyl ether (30 ml) was slowly added and the mixture was stirred for 30 minutes. The solid was isolated by filtration, washed with diethyl ether, and dried to give 0.78 g of (3R)-3-benzyl-1-methanesulfonylpiperazine hydrochloride. M.p. 204-208°C.

¹H-NMR (DMSO-d₆): δ 2.87-3.25 (m, 8H); 3.35-3.65 (m, 4H); 7.25-7.40 (m, 5H).

N-[(1R)-2-((2R)-2-Benzyl-4-methanesulfonylpiperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxoethyl]-N-methylcarbamic acid tert-butyl ester.




(2R)-((tert-Butoxycarbonylmethylamino)-3-(2-naphthyl)propionic acid (0.86 g; 2.6

mmol) was dissolved into dichloromethane (25 ml) and placed under an atmosphere of nitrogen. (3R)-3-Benzyl-1-methanesulfonylpiperazine hydrochloride (0.75 g; 2.6 mmol), N-methylmorpholine (0.26 g; 2.6 mmol), 1-hydroxy-7-azabenzotriazole (0.39 g; 2.9 mmol) and N-(3-dimethylaminopropyl)-
 5 N'-ethylcarbodiimide hydrochloride (0.55; 2.9 mmol) were added and the mixture was stirred overnight at ambient temperature. Dichloromethane (25 ml) was added and the mixture was washed with water (3 x 15 ml) and dried over magnesium sulfate. The solvent was evaporated in vacuo to give 1.3 g of an oily residue which was chromatographed on silica gel using a mixture of heptane
 10 and ethyl acetate (1:1) as eluent. This afforded 0.65 g of N-[(1R)-2-((2R)-2-benzyl-4-methanesulfonylpiperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxo-ethyl]-N-methylcarbamic acid tert-butyl ester as a solid.

M.p. 76-80°C.

15


(2R)-1-((2R)-2-Benzyl-4-methanesulfonyl-piperazin-1-yl)-2-methylamino-3-(naphth-2-yl)-propan-1-one.



N-[(1R)-2-((2R)-2-Benzyl-4-methanesulfonylpiperazin-1-yl)-(1R)-1-(naphth-2-ylmethyl)-2-oxo-ethyl]-N-methylcarbamic acid tert-butyl ester (0.63 g; 1.1 mmol)
 20 was dissolved into dichloromethane (10 ml) with stirring and placed under an atmosphere of nitrogen. Trifluoroacetic acid (10 ml) was added and the mixture was stirred for 15 minutes. The reaction mixture was diluted with dichloromethane (75 ml) and placed on an ice-bath. Saturated aqueous sodium
 25 bicarbonate solution was added and then an aqueous potassium carbonate solution until pH 10. The phases were separated and the aqueous phase was extracted with dichloromethane (50 ml). The combined organic extracts were dried over magnesium sulfate and the solvent was evaporated in vacuo. This
 30 afforded 0.51 g of (2R)-1-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-2-methylamino-3-(naphth-2-yl)propan-1-one as a solid.

M.p. 165-168°C.

5 ((3E)-4-(N-((1R)-2-((2R)-2-Benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-yl-methyl)-2-oxo-ethyl)-N-methylcarbamoyl)-1,1-dimethyl-but-3-enyl)carbamic acid tert-butyl ester.



10 (2E)-5-(tert-Butyloxycarbonylamino)-5-methylhex-2-enoic acid (0.49 g; 2.0 mmol) was dissolved into dichloromethane (10 ml) and placed under an atmosphere of nitrogen. 1-Hydroxy-7-azabenzotriazole (2.0 mmol) and N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (2.0 mmol) were added and the mixture was stirred for 15 minutes. A solution of 1-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-(2R)-2-methylamino-3-naphth-2-yl-propan-1-one (0.51 g; 1.1 mmol) in dichloromethane (10 ml) was added and the mixture was stirred overnight. Dichloromethane (50 ml) was added and the mixture was washed with water (2 x 50 ml). The combined aqueous phases were extracted with dichloromethane (50 ml). The combined organic extracts were dried over magnesium sulfate and the solvent was evaporated in vacuo. This afforded 1.0 g residue which was chromatographed on silica gel using a mixture of heptane and ethyl acetate (3:7) as eluent. This afforded 0.55 g of

15 ((3E)-4-(N-((1R)-2-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-yl-methyl)-2-oxoethyl)-N-methylcarbamoyl)-1,1-dimethylbut-3-enyl)carbamic acid tert-butyl ester.

25 TLC; R_f = 0.33 (heptane/ethyl acetate = 3:7).

((3E)-4-(N-((1R)-2-((2R)-2-Benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxoethyl)-N-methyl-carbamoyl)-1,1-dimethylbut-3-enyl)carbamic acid tert-butyl ester (0.54 g; 0.78 mmol) was dissolved in dry dichloromethane (10 ml) with stirring and trifluoroacetic acid (10 ml) was added. The mixture was

30


stirred for 8 min. at ambient temperature and then diluted with dichloromethane (50 ml). The mixture was placed on an ice-bath and saturated aqueous sodium bicarbonate solution was added and then an aqueous potassium carbonate solution until pH 10. The phases were separated and the aqueous phase was
5 extracted with dichloromethane (20 ml). The combined organic extracts were dried over magnesium sulfate and the solvent was evaporated in vacuo. This afforded 0.33 g of (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxoethyl)-N-methylamide.

10

¹H-NMR (CDCl₃): δ 1.17 (s, 3H); 1.20 (s, 3H); 1.8-3.65 (complex m); 4.10 (d, 1H); 4.55 + 4.88 (d+m, 1H); 5.82 + 6.03 (t+dd, 1H); 6.28 (dd, 1H); 6.81 + 6.95 (dt+dt, 1H); 7.15-7.80 (m, 12H).

Claims:

1. A compound of general formula I



formula I

wherein

10 R¹ is hydrogen, or C₁₋₈-alkyl optionally substituted with aryl;

a and b are independently 1 or 2;

c and d are independently 0, 1, or 2;

15


c + d is 0, 1, or 2;

20 R², R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹⁰, and R¹¹ are independently hydrogen, aryl optionally substituted with halogen, amino, C₃₋₆-cycloalkyl, hydroxy, aryl, -COOR²², or -CONR²³R²⁴, or C₁₋₆-alkyl optionally substituted with halogen, amino, C₃₋₆-cycloalkyl, hydroxy, aryl, -COOR²², or -CONR²³R²⁴;


R³ and R⁴ can be taken together to form =O or =S;

25 R⁸ and R⁹ can be taken together to form =O or =S;

R⁵ is hydrogen,



or C₁₋₆-alkyl optionally substituted with aryl, hydroxy, C₃₋₆-cycloalkyl, amino, -





5 R, R' and R'' are independently hydrogen or C₁₋₆-alkyl;

R³⁰ and R³¹ are C₁₋₆-alkyl optionally substituted with aryl, hydroxy, C₃₋₆-cycloalkyl, or amino;

10 R²², R²³, R²⁴, R²⁵, R²⁶, and R²⁷ are independently hydrogen or C₁₋₆-alkyl;



when R³ and R⁴ are taken together to form =O or =S, E is



15

when R³ or R⁴ is hydrogen, aryl optionally substituted with halogen, amino, C₃₋₆-cycloalkyl, hydroxy, aryl, -COOR²², or -CONR²³R²⁴, or C₁₋₆-alkyl optionally substituted with halogen, amino, C₃₋₆-cycloalkyl, hydroxy, aryl, -COOR²², or

20 -CONR²³R²⁴, E is



R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

5

any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;

n , m and q are independently 0, 1, 2, or 3;

10

o and p are independently 0 or 1;

M is $-CH=CR^{21}-$, $-O-$, $-S-$, or a valence bond ;


15

Q is $-CH=CR^{21}-$, $-O-$, or $-S-$;

R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

20


G is hydrogen,



wherein R^{28} is hydrogen, halogen, C_{1-6} -alkyl, C_{1-6} -alkoxy or aryl;

5 r is 0, 1, or 2;

J is hydrogen,




10 wherein R^{29} is hydrogen, halogen, C_{1-6} -alkyl, C_{1-6} -alkoxy or aryl;

t is 0, 1, or 2;

or a pharmaceutically acceptable salt thereof.

15

2. A compound of the general formula II




formula II


5

wherein

E is



10



R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

15 any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;

n , m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;


M is $-\text{CH}=\text{CR}^{21}-$, $-\text{O}-$, $-\text{S}-$, or a valence bond ;

- 5 R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl; and

$\text{R}^1, \text{R}^2, \text{R}^5, \text{R}^6, \text{R}^7, \text{R}^8, \text{R}^9, \text{R}^{10}, \text{R}^{11}, \text{G}, \text{J}, \text{a}, \text{b}, \text{c},$ and d are defined as in claim 1;


- 10 or a pharmaceutically acceptable salt thereof.

3. The compound of the general formula III




formula III

wherein E is



20



R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;

n , m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;

M is $-\text{CH}=\text{CR}^{21}-$, $-\text{O}-$, $-\text{S}-$, or a valence bond ;


R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

R^8 and R^9 are independently hydrogen, aryl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-\text{COOR}^{22}$, or $-\text{CONR}^{23}\text{R}^{24}$, or C_{1-6} -alkyl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-\text{COOR}^{22}$, or $-\text{CONR}^{23}\text{R}^{24}$; and

G , J , R^1 , R^2 , R^5 , R^6 , R^7 , R^{22} , R^{23} , R^{24} , a , and b are as defined in claim 1;

or a pharmaceutically acceptable salt thereof.

4. The compound according to any one of the claims 2-3 wherein R^5 is hydrogen or C_{1-6} -alkyl optionally substituted with aryl, hydroxy, C_{3-6} -cycloalkyl, amino, -



R , R' and R'' are independently hydrogen or C_{1-6} -alkyl;

R^{25} , R^{26} , and R^{27} are independently hydrogen or C_{1-6} -alkyl; and


$R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, E, G, J, a, b, c,$ and d are as defined in any of the preceeding claims;

or a pharmaceutically acceptable salt thereof.


5

5. The compound according to any one of the claims 2-4, wherein

E is



10



$R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R^{17}, R^{18}, R^{19},$ and R^{20} are independently hydrogen or methyl;
 n, m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;

15

M is $-CH=CR^{21}-$, $-O-$, or a valence bond ;

R^{21} is hydrogen, methyl or benzyl; and

20

$R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, G, J, a, b, c,$ and d are as defined in any of the preceeding claims;

or a pharmaceutically acceptable salt thereof.



25

6. The compound according to claim 1 wherein

R^3 and R^4 are independently hydrogen, aryl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$, or C_{1-6} -alkyl optionally substituted with halogen, amino, C_{3-6} -cycloalkyl, hydroxy, aryl, $-COOR^{22}$, or $-CONR^{23}R^{24}$;

R^{22} , R^{23} , and R^{24} are independently hydrogen or C_{1-6} -alkyl;

10 E is



R^{12} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

15

any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;

n , m and q are independently 0, 1, 2, or 3;

20

o and p are independently 0 or 1;

Q is $-CH=CR^{21}-$, $-O-$, or $-S-$;


R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl; and

$R^1, R^2, R^5, R^6, R^7, R^8, R^9, R^{10}, R^{11}, G, J, a, b, c,$ and d are as defined in claim 1

5

or a pharmaceutically acceptable salt thereof.



7. A compound of the general formula IV



formula IV

10

wherein E is



15 $R^{12}, R^{17}, R^{18}, R^{19},$ and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

any two of $R^{17}, R^{18},$ and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;

20

n, m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;


5 Q is $-\text{CH}=\text{CR}^{21}-$, $-\text{O}-$, or $-\text{S}-$;

R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl; and

10 R^1 , R^5 , R^6 , G, J, a, and b are as defined in claim 1;

or a pharmaceutically acceptable salt thereof.



15 8. A compound of the general formula V



formula V

wherein E is

20



R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , and R^{20} are independently hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

any two of R^{17} , R^{18} , and R^{19} can independently be joined together to form an C_{1-6} -alkylene bridge;


n , m and q are independently 0, 1, 2, or 3;

o and p are independently 0 or 1;

M is $-CH=CR^{21}-$, $-O-$, $-S-$, or a valence bond ;

R^{21} is hydrogen or C_{1-6} -alkyl optionally substituted with halogen, amino, hydroxyl or aryl;

R^5 is hydrogen or C_{1-6} -alkyl optionally substituted with aryl, hydroxy, C_{3-6} -cycloalkyl,

amino, $-COOR^{25}$, $-CONR^{26}R^{27}$, $-NR'R''$,  ;

R , R' and R'' are independently hydrogen or C_{1-6} -alkyl;

R^{25} , R^{26} , and R^{27} are independently hydrogen or C_{1-6} -alkyl; and

G , J , a , b , R^1 , and R^6 are as defined in claim 1;

or a pharmaceutically acceptable salt thereof.

5 9. The compound according to any one of the claims 1-8 wherein

R¹ is hydrogen or methyl;

or a pharmaceutically acceptable salt thereof.

10

10. The compound according to claims 1-8 or 9 selected from the group consisting of

15 (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

(2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-(cyclopropylmethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

20 (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-(hydroxymethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

(2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxo-1,4-diazepane-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

25

piperidine-4-carboxylic acid N-methyl-N-((1R)-2-(2-naphthyl)-1-((2R)-2-((2-naphthyl)methyl)-3-oxopiperazine-1-carbonyl)ethyl)amide,

30 (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-(dimethylcarbamoylmethyl)-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

(2E)-5-methyl-5-(methylamino)hex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,

35

- (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R,6S)-2-benzyl-6-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,
- 5 (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(biphenyl-4-yl)ethyl)-N-methylamide,
- (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-methanesulfonylpiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,
- 10 (2E)-5-amino-5-methylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-4-methanesulfonylpiperazine-1-carbonyl)-2-benzyloxyethyl)-N-methylamide,
- (2E)-5-Amino-3,5-dimethylhex-2-enoic acid N-((1R)-1-((2R)-2-benzyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,
- 15 (2E)-5-Methyl-5-methylaminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,
- (2E)-5-Amino-5-methylhex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide,
- 20 (2E)-5-Methyl-5-aminohex-2-enoic acid N-((1R)-1-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazine-1-carbonyl)-2-(2-naphthyl)ethyl)-N-methylamide,
- (2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-methyl-N-((1R)-2-(2-naphthyl)-1-((2R)-3-oxo-2-((2-thienyl)methyl)piperazine-1-carbonyl)ethyl)amide,
- 25 (2E)-4-(1-Aminocyclobutyl)but-2-enoic acid N-((1R)-2-((2R,5R)-2-benzyl-5-methyl-3-oxopiperazin-1-yl)-1-((2-naphthyl)methyl)-2-oxoethyl)-N-methylamide,
- 30 (2E)-5-Methyl-5-methylamino-hex-2-enoic acid N-methyl-N-((1R)-1-(2-naphthyl)methyl-2-oxo-2-((2R)-3-oxo-2-((2-thienyl)methyl)piperazin-1-yl)ethyl)amide,

(2E)-5-Amino-5-methyl-hex-2-enoic acid N-((1R)-2-((2R)-2-benzyl-4-methanesulfonyl-piperazin-1-yl)-1-(naphth-2-ylmethyl)-2-oxo-ethyl)-N-methylamide,

or a pharmaceutically acceptable salt thereof.

5

11. A pharmaceutical composition comprising, as an active ingredient, a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

10

12. The composition according to claim 11 in unit dosage form, comprising from about 10 to about 200 mg of the compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof.

15

13. A pharmaceutical composition for stimulating the release of growth hormone from the pituitary, the composition comprising, as an active ingredient, a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

20

14. A pharmaceutical composition for administration to animals to increase their rate and extent of growth, to increase their milk and wool production, or for the treatment of ailments, the composition comprising, as an active ingredient, a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

25

15. A pharmaceutical composition according to any one of the claims 11-14 for oral, nasal, transdermal, pulmonal, or parenteral administration.

30

16. A method of stimulating the release of growth hormone from the pituitary, the method comprising administering to a subject in need thereof an effective amount of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof or of a composition according to any one of the claims 11-

35

13 or 15.

17. A method according to claim 16, wherein the effective amount of the compound according to any one of the claims 1-10 or pharmaceutically acceptable salt or ester thereof is in the range of from about 0.0001 to about 100
5 mg/kg body weight per day, preferably from about 0.001 to about 50 mg/kg body weight per day.
18. A method for increasing the rate and extent of growth of animals to increase their milk and wool production, or for the treatment of ailments, the method
10 comprising administering to a subject in need thereof an effective amount of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof or of a composition according to any one of the claims 11, 12, 14 or 15.
- 15 19. The method according to any one of the claims 16-18, wherein said administration is carried out by the oral, nasal, transdermal, pulmonal, or parenteral route.
- 20 20. Use of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof for the preparation of a medicament.
21. Use of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for stimulating the release of growth hormone from the pituitary.
25
22. Use of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for administration to animals to increase their rate and extent of growth, to increase their milk and wool production, or for the treatment of ailments.
30
23. Use of a compound according to any one of the claims 1-10 or a pharmaceutically acceptable salt thereof for the preparation of a medicament for the treatment of stimulation of growth hormone release in the elderly; prevention of catabolic side effects of glucocorticoids, prevention and treatment of
35 osteoporosis, stimulation of the immune system, acceleration of wound healing,

- accelerating bone fracture repair, treatment of growth retardation, treating renal failure or insufficiency resulting from growth retardation, treatment of physiological short stature including growth hormone deficient children and short stature associated with chronic illness, treatment of obesity and growth retardation
- 5 associated with obesity, treating growth retardation associated with the Prader-Willi syndrome and Turner's syndrome; accelerating the recovery and reducing hospitalization of burn patients; treatment of intrauterine growth retardation, skeletal dysplasia, hypercortisolism and Cushing's syndrome; induction of pulsatile growth hormone release; replacement of growth hormone in
- 10 stressed patients, treatment of osteochondrodysplasias, Noonan's syndrome, schizophrenia, depressions, Alzheimer's disease, delayed wound healing and psychosocial deprivation, treatment of pulmonary dysfunction and ventilator dependency, attenuation of protein catabolic responses after major surgery, reducing cachexia and protein loss due to chronic illness such as cancer or AIDS;
- 15 treatment of hyperinsulinemia including nesidioblastosis, adjuvant treatment for ovulation induction; to stimulate thymic development and prevent the age-related decline of thymic function, treatment of immunosuppressed patients, improvement in muscle strength, mobility, maintenance of skin thickness, metabolic homeostasis, renal homeostasis in the frail elderly, stimulation of osteoblasts,
- 20 bone remodelling and cartilage growth, stimulation of the immune system in companion animals and treatment of disorder of aging in companion animals, growth promoter in livestock and stimulation of wool growth in sheep.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 97/00186

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: C07D 241/08, C07D 241/04, C07D 243/08, C07D 401/12, C07D 409/06,
A61K 31/495

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: C07D, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CASONLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 9517423 A1 (NOVO NORDISK A/S), 29 June 1995 (29.06.95) --	1-23
A	Chem. Pharm. Bull., Volume 44, No 4, 1996, Tetsushi Yamashita et al, "Structure-Activity Relationships of Dermorphin Analogues Containing Chiral Piperazin-2-one and Piperazine Derivatives" page 856 - page 859 -- -----	1-23

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 August 1997

Date of mailing of the international search report

14.08.97

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK97/00186

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-19
because they relate to subject matter not required to be searched by this Authority, namely:
See PCT Rule 39.1.(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☒ Claims Nos.: 23
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claim 23 is not clear and concise, as it is not likely that all these medical disorders (such as, Schizophrenia, depression, Alzheimer, pulmonary dysfunction) are related to growth hormone.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

01/07/97

International application No.

PCT/DK 97/00186

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9517423 A1	29/06/95	AU 1272495 A	10/07/95
		CA 2179597 A	29/06/95
		CN 1138335 A	18/12/96
		CZ 9601834 A	12/02/97
		EP 0736039 A	09/10/96
		FI 962584 A	20/06/96
		HU 73497 A	28/08/96
		HU 9501947 D	00/00/00
		IL 112112 D	00/00/00
		NO 962665 A	23/08/96
		NZ 277486 A	24/03/97
		PL 315113 A	14/10/96
		SK 82096 A	08/01/97
		ZA 9410261 A	23/06/95

THIS PAGE BLANK (USPTO)